

Russian Initiative – Central Asia Rangeland Activity

Draft Version

METHODOLOGY

**For Assessing Rangeland Vegetation
in the Action Sites (*Karakalpakstan & Tajikistan*)**



Prepared by Mounir Louhaichi, Toshpulat Rajabov and Makhmud Shaumarov

05 July 2015

Russian Initiative – Central Asia Rangeland Activity

Objectives of the research

The research objectives has two folds aiming at:







1. Fine tuning/developing a toolkit for monitoring and assessing rangeland vegetation at the landscape level using up-to-date technologies; and
2. Carrying out an integrated assessment (characterizing/mapping) of grazing driven vegetation dynamics in two selected and distinctive ecological zones (desert versus mountain).

Field experimental design




We selected 2 sites which are distinctive in terms of environmental condition (climate, soil, topography and vegetation structure). The first action site represents typical sandy rangelands of Kyzylkum desert in Karakalpakstan, whereas the second site is located in Kurama mountain ranges in North Tajikistan. The current rangeland vegetation in each site represent the dominated type which is highly shaped by livestock grazing. The dominant native species in the Kyzylkum desert in Karakalpakstan is white haloxylon (*Haloxylon persicum*) while the invasive species *Ferula assa-foetida* L. Represents a major threat to the entire ecosystem.



Characteristics of dominant species

Growth of <i>Ferula assa-foetida</i> L.		
		
Seedlings of <i>Ferula</i> at an early stage		
		
Mature plant with green leaves	Mature plant with yellow leaves	Dead <i>Ferula</i> (> 8 years) old


Russian Initiative – Central Asia Rangeland Activity

Dominant native plant species		
		
White haloxyton	Artemisia	Carex (annual)

Sandy desert rangelands



We used a grazing gradient approach as a main tool to detect fine-scale changes of vegetation composition and its structure. Three monitoring sites with different level of livestock grazing were selected for conducting the vegetation surveys: rangeland areas around 2 watering wells and one rangeland area with no livestock grazing. The rangeland vegetation around 2 watering wells is surveyed as a distance away from the watering well. Three transects radiating in 3 directions from the center of the well were allocated at each 120 degrees apart from North. The length of each transects ranges between 3.0-3.5 km from the center of the well. Along each transect as a distance away from the well 3 stops were selected to detect vegetation changes caused by different level of livestock grazing. In total 9 stops were monitored around each watering well.



Mountain rangelands




In this action site, we have selected North versus South facing slopes since the aspect (denotes the compass direction in which the slope of a mountain faces) has great influence on long term vegetation formation and its current condition. Selected North versus South facing slopes are also characterized by different level of grazing intensity.

Russian Initiative – Central Asia Rangeland Activity



The slopes with different grazing intensity were separately selected in winter and summer pastures. Vegetation surveys were done on North and South aspects in each selected V-shape slope as a distance away from the livestock flock (up to 2.5-3.0 km). In addition, the vegetation of other types of landscapes as gently rolling flat hills in winter pastures, rocky mountains and flat rangelands in a proximity of mountains were covered during vegetation surveys.



Sampling of plant community data

Vegetation data of spring season were collected during 01-05 May in Karakalpakstan and 15-20 May in Tajikistan. The vegetation measurements basically included biomass, cover and density of perennial plants, biomass and density of annuals.

Description of the plant community was done using 50 x 2 m quadrat in scarce vegetation of sand rangelands whereas mountain vegetation was described using 10 x 2 m quadrat due to its high species richness.

Russian Initiative – Central Asia Rangeland Activity



The total numbers of shrubs of each species were counted and separated into 3 size categories (big, medium, small). For each species within every size category, 3 representative plants were clipped for determination of annual green biomass using the reference technique (Ref).



The cover of shrub species was determined along a line intercept as shown below.



Biomass production of ephemerals and ephemeroïds was identified within 1 x 1 m frame quadrates, randomly distributed with 3 replications.




Russian Initiative – Central Asia Rangeland Activity



At the same time we used the Digital Vegetation Charting Technique (DVCT) for total cover and density assessment of vegetation cover (Ref). In addition data collected with DVCT will be used also as a training site for supervised classification of satellite RS.




Density estimation of Carex (annual species) using DVCT



Russian Initiative – Central Asia Rangeland Activity

Canopy cover estimation using DVCT



Developing allometric equation to estimate Artemisia biomass using cover, height and diameter as surrogate



Linking near earth RS to satellite RS (Chandra to add)

Perspective

The steps described above have to be repeated again during the fall to capture the seasonal variations (temporal resolution). This frequency is needed to document the status of rangeland vegetation (productivity and quality) before and after grazing.

The obtained ground truth vegetation data and key findings in both action sites will be overlaid to environmental and management conditions. The results then will be incorporated into GIS and RS technologies to characterize spatial and temporal dynamics of rangeland vegetation. Satellite remote sensing data will be incorporated at different spectral and spatial resolutions. The most cost effective approach and tools will be outscaled. The results may lay on the basis of development of operative methods in assessing rangeland condition of different ecological zones.

Note: The dense vegetation in the sandy rangelands of Kyzylkum desert in Karakalpakstan makes difficult for coarse RS data to assess rangeland value (species composition, palatability, etc.). The site is invaded by *Ferula*. Although the species is unpalatable, *Ferula assa-foetida* L. is used as spice, as aphrodisiac and as herbal remedy. It was used since ancient times. It is an oleo gum resin which exudes out of the rhizomes. Normally it is used as an antifatulent, digestive aid. Pharmacologically used as antimicrobial, antiasthmatic, antiepileptic and also reported to have contraceptive/abortifacient activity.

The strong smell is inherent and characteristic of the oleogum resin. It has a very strong unpleasant smell.

Russian Initiative – Central Asia Rangeland Activity

