## OPTIONS méditerranéennes

# Efficiency and resilience of forage resources and small ruminant production to cope with global challenges in Mediterranean areas

Edited by: A. López-Francos, M. Jouven, C. Porqueddu, H. Ben Salem, A. Keli, A. Araba, M. Chentouf





**SERIES A: Mediterranean Seminars** 2021 – Number 125





### **CIHEAM**

#### Centre International de Hautes Etudes Agronomiques Méditerranéennes

#### International Centre for **Advanced Mediterranean Agronomic Studies**

Président / President: Mohammed SADIKI Secretariat General / General Secretariat: Plácido PLAZA

11, rue Newton, 75116 Paris, France Tél.: +33 (0) 1 53 23 91 00 - Fax: +33 (0) 1 53 23 91 01 / 02 secretariat@ciheam.org www.ciheam.org

Le Centre International de Hautes Etudes Agronomiques Méditerranéennes (CIHEAM) a été créé, à l'initiative conjointe de l'OCDE et du Conseil de l'Europe, le 21 mai 1962. C'est une organisation intergouvernementale qui réunit aujourd'huitreize Etats membres du bassin méditerranéen (Albanie, Algérie, Egypte, Espagne, France, Grèce, Italie, Liban, Malte, Maroc, Portugal, Tunisie et Turquie).

Le CIHEAM se structure autour d'un Secrétariat général situé à Paris et de quatre Instituts Agronomiques Méditerranéens (IAM), localisés à Bari (Italie), Chania (Grèce), Montpellier (France) et Saragosse (Espagne).

Avec au cœur de son action trois missions fondamentales (formation, recherche, coopération), le CIHEAM s'est progressivement imposé comme une référence dans ses domaines d'activité : l'agriculture, l'alimentation et le développement rural durable en Méditerranée.

Founded in 1962 at the joint initiative of the OECD and the Council of Europe, the International Centre for Advanced Mediterranean Agronomic Studies (CIHEAM) is an intergovernmental organisation comprising thirteen member countries from the Mediterranean Basin (Albania, Algeria, Egypt, Spain, France, Greece, Italy, Lebanon, Malta, Morocco, Portugal Tunisia and Turkey).

CIHEAM is made up of a General Secretariat based in Paris and four Mediterranean Agronomic Institutes (MAI) located in Bari (Italy), Chania (Greece), Montpellier (France) and Zaragoza (Spain).

In pursuing its three central missions (education, research and cooperation) CIHEAM has established itself as a reference in its fields of activity: Mediterranean agriculture, food and sustainable rural development.

#### **Instituts Agronomiques Méditerranéens Mediterranean Agronomic Institutes**

#### Bari - Chania - Montpellier - Zaragoza

#### IAM-Bari

Dir.: Maurizio RAELI Via Ceglie, 9

70010 Valenzano, Bari, Italy

Tel.: (+39) (080) 4606 111 - Fax: (+39) (080) 4606 206

iamdir@iamb.it www.iamb.it

#### IAM-Montpellier

Dir.: Pascal BERGERET 3191. Route de Mende

34093 Montpellier Cedex 5, France

Tel.: (+33) (0)4 67 04 60 00 - Fax: (+33) (0)4 67 54 25 27 Tel.: (+34) 976 716000 - Fax: (+34) 976 716001

pascal.bergeret@iamm.fr et/and sciuto@iamm.fr

www.iamm.fr

#### IAM-Chania

Dir.: Giorgios BAOURAKIS

P.O. Box 85

73100 Chania, Crete, Greece

Tel.: (+30) 28210 35000 - Fax: (+30) 28210 35001

baouraki@maich.gr www.maich.gr

#### IAM-Zaragoza

Dir.: Raúl COMPES Avda. Montañana, 1005 50059 Zaragoza, Spain

iamz@iamz.ciheam.org www.iamz.ciheam.org

### Efficiency and resilience of forage resources and small ruminant production to cope with global challenges in Mediterranean areas

Editors: A. López-Francos, M. Jouven, C. Porqueddu, H. Ben Salem, A. Keli, A. Araba, M. Chentouf

Proceedings of the 1st Joint Meeting of the FAO-CIHEAM Network for Research and Development in Sheep and Goats (Subnetworks on Nutrition and Production Systems) and the FAO-CIHEAM Subnetwork on Mediterranean Forage Resources, jointly organised by the École Nationale d'Agriculture de Meknès (ENA), under the aegis of the Moroccan Ministry of Agriculture, Fisheries, Rural Development, Water and Forests (MAPMDREF), and by the Mediterranean Agronomic Institute of Zaragoza/International Centre of Advanced Mediterranean Agronomic Studies (CIHEAM-Zaragoza); with the collaboration of the Food and Agriculture Organization of the United Nations (FAO) and the H2020 Project iSAGE (Innovation for Sustainable Sheep and Goat Production in Europe, GA no. 679302).

Meknes (Morocco), 23 to 25 October 2019







With the collaboration of:





## OPTIONS méditerranéennes

Head of publication: Plácido Plaza

2021 Series A: Mediterranean Seminars

Number 125



Centre International de Hautes Etudes Agronomiques Méditerranéennes International Centre for Advanced Mediterranean Agronomic Studies L'édition technique, la maquette et la mise en page de ce numéro d'Options Méditerranéennes ont été réalisées par INO Reproducciones, S. A. Technical editing, layout and formatting of this edition of Options Méditerranéennes was carried out by INO Reproducciones, S. A.

> Crédits des photos de couverture / Cover photo credits: Abdelhafid Keli, Antonio López-Francos

> > Tirage / Copy number Printer: INO Reproducciones, S.A. Pol. Malpica, calle E, 32-39 (INBISA II, Nave 35) 50016 Zaragoza-Spain Dep. Legal: Z-2893-91

#### Fiche bibliographique / Cataloguing data:

Efficiency and resilience of forage resources and small ruminant production to cope with global challenges in Mediterranean areas. A. López-Francos, M. Jouven, C. Porqueddu, H. Ben Salem, A. Keli, A. Araba, M. Chentouf (eds). Zaragoza: CIHEAM/ENA 2021, 704 p. (Options Méditerranéennes, Series A: Mediterranean Seminars. No. 125)

Catalogue des numéros d'Options Méditerranéennes sur /
Catalogue of Options Méditerranéennes issues on:
www.ciheam.org/publications

ISSN: 1016-121-X – ISBN: 978-2-85352-608-1 © CIHEAM, 2021

Reproduction partielle ou totale interdite sans l'autorisation du CIHEAM

Reproduction in whole or in part is not permitted without the consent of CIHEAM

## Developing a toolbox for rangeland restoration/rehabilitation in arid environments

M. Louhaichi<sup>1,\*</sup>, P.M. Moyo<sup>1</sup>, S. Hassan<sup>1</sup> and A. Ouled Belgacem<sup>2</sup>

<sup>1</sup>International Center for Agricultural Research in the Dry Areas, Amman (Jordan)

<sup>2</sup>International Center for Agricultural Research in the Dry Areas, Dubai (UAE)

\*e-mail: m.louhaichi@cgiar.org

Abstract. Rangelands are recognized for their importance and value in providing society with valuable products and ecosystem services. In such ecosystems, effective management is needed for sustainable plant growth and survival in a context characterized by rainfall unreliability, poor soil nutrient status and high uncontrolled grazing. Therefore, cost-effective techniques/tools for slowing down and eventually reversing this degradation are needed. This paper promotes the identification and association of various tools for degraded arid ecosystems as strategies aimed at rangeland restoration/ rehabilitation. These strategies are founded on science-based evidence and experienced practitioners. For arid rangelands, the preference of applying an isolated rehabilitation technique may not halt degradation since the heterogeneity of the geomorphology, even at micro scale, and the landscape requires very often the use of different tools. For instance, direct seeding in degraded rangelands needs to be combined with soil scarification, when the crust dominates the soil surface, to reactivate the soil water retention and increase seedling emergence and germination possibilities. To achieve this goal, rehabilitation options identified for a specific degraded rangeland need to be holistically integrated with land degradation indicators in a manual-style decision support system for the long-term sustainable production of rangelands in arid environments.

**Keywords.** Dry areas – Land degradation – Sustainable rangeland management – Restoration.

#### Développer une boîte à outils pour la restauration / réhabilitation des parcours en milieu aride

Résumé. Les parcours jouent un rôle écologique et économique très important et fournissent aux sociétés pastorales des produits et des services écosystémiques précieux. Dans de tels écosystèmes, une gestion efficace est nécessaire pour la croissance et la survie durables des plantes dans un contexte caractérisé par une aridité climatique et édaphique très accentué et soumis à une pression de pâturage élevée et non contrôlée. Par conséquent, il devient nécessaire de développer des techniques / outils rentables pour ralentir et éventuellement inverser cette dégradation. Cet article vise l'identification et la combinaison de divers outils pour les écosystèmes arides dégradés en tant que stratégies en vue de restaurer/réhabiliter les parcours naturels. Ces stratégies sont fondées sur des preuves scientifiques et des pratiques validées. Pour les parcours arides, le recours préférentiel à une technologie de réhabilitation isolée ne peut pas arrêter la dégradation car l'hétérogénéité de l'écosystème, même à petite échelle, nécessite très souvent la combinaison de plusieurs outils différents. Par exemple, la plantation d'arbustes fourragers sur des pentes relativement raides doit être associée à une technique de collection des eaux pluviales, pour tenir en compte du caractère torrentiel des en milieu aride. Pour atteindre cet objectif, les options de réhabilitation identifiées pour un parcours dégradé doivent être intégrées de manière holistique aux indicateurs de dégradation des terres dans un système d'aide à la décision sous la forme d'un référentiel technique pour assurer une gestion durable de parcours dans en milieux arides.

Mots-clés. Zones arides – Dégradation du sol – Gestion durable des parcours – Restauration.

#### I - Introduction

The arid and semi-arid areas are estimated to cover 35% of the earth's land surface and are dominated by rangelands. They are major providers of critical ecosystem goods and services, including food, water, and livelihood sources for many of the world's poor (Milder et al., 2010). Unfortu-

nately, these resources are on steady decline. For many decades national programs and development projects have attempted to reverse the trend of degradation, but in most cases the results were discouraging. Degradation which took place over hundreds of years cannot be reversed using quick fix solutions without considering the underlying causes. In fact, degradation is a result of poor management strategies interacting with other factors, such as climate change, which have continually increased the pressure on natural resources (Ouled Belgacem and Louhaichi, 2013). It is crucial to understand the causes of rangeland degradation and find solutions mobilizing a suite of available tools in a community-based participatory approach.

In this context, this toolbox has been developed to address rangeland rehabilitation at the landscape-level, taking into consideration the biophysical and socioeconomic linkages and trade-offs existing between the different land uses.

#### II - Sustainable Rangeland Management Practices

There are several sustainable rangeland management practices (SRMPs) that have been developed over the years. These SRMPs combine both indigenous knowledge and science-based interventions. Due to limited space (page number), in this paper we will only focus on key practices and approaches needed to ensure sustainability of rangeland restoration and rehabilitation.

#### 1. Participatory rangeland governance

Sustainable rehabilitation of degraded rangelands that are dominated by collective and/or tribal ownership is a challenging task. The institutional arrangement to tackle this complex issue has been rather sectorial and fragmented. Earlier, the 'top–down' approach, which emphasizes technical solutions and neglects the social context, was the more common form of intervention. Therefore, in response to frequent failures of the top-down approach, efforts were deployed towards 'participatory development' as an alternative approach. National governments have been slow to adopt this participatory approach, but recent experiences suggest that integrated and participatory rangeland governance approaches may lead to more sustainable resource management. Rangeland governance can be defined as "local people's participation in managing the rangeland resources that they themselves use, in a sustainable manner" (Rist *et al.*, 2007). Such arrangement of development aims to organize people on a decentralized basis and to apply participatory tools in order to effectively empower local people (Nefzaoui *et al.*, 2014).

#### 2. Water harvesting techniques

Water is an important resource in arid and semi-arid rangelands, but often it is wasted or allowed to erode the landscape. One way to address this is through simple, cost-effective water harvesting techniques (WHT). This practice has been used for thousands of years in arid and semi-arid regions of the world to supplement scarce water resources. In general, the interventions are used to increase soil moisture content, vegetation cover, and productivity. WHT can also control soil erosion and reduce the impact of drought. Experience of the last two decades provides increasing evidence that WHT can mitigate the increasing variability of rainfall. Among the widely used micro catchment WH techniques are contour ridges, semicircular and trapezoidal bunds and small runoff basins. When implementing WHT, field conditions such as slope of the terrain, soil properties, land use and land cover need to be assessed before choosing and constructing water harvesting structures. For large-scale implementation, a special plough can be designed to construct intermittent and continuous contour ridging (Oweis, 2016). In small areas and depending on the land characteristics, different manual constructing bunds can be implanted, includeing semi-circular bunds, stone bunds, earth basins and *Maskat*.

#### 3. Soil surface scarification

Soil surface scarification is a technique that breaks up compacted surface soil to improve water infiltration. The main purposes of this practice are to facilitate germination and emergence of soil seeds and to create a favorable seedbed. This practice is commonly used to ensure a successful regeneration of the natural vegetation, either by sowing or by natural rehabilitation. It is recommended to use a chisel plow or a pitting machine to disturb the top crusty soil (upper 5-10 cm). This intervention should take place 1 to 2 weeks before the expected rainfall events (Gauthier, 2016).

#### 4. Reseeding

Reseeding can be described as the process of introducing seeds of a new or fresh crop to replenish a depleted soil seed bank. Low-productive rangelands, which result in a lack of balance in forage production, must be rehabilitated by inexpensive methods such as reseeding to provide more and better-quality forage to support livestock and protect the soil. Reseeding has a potential of yielding high plant density at low costs and it is a cheap means for providing an adequate feed for livestock and a greater income from the grazing operation. Reseeding is appropriate for moderate to highly-degraded rangelands, where average annual rainfall is above 350 mm. It should be done only as part of a good rangeland management plan to improve both the reseeded area and the native range. Reseeding is necessary to increase nutritional value of forage, fill in bare spots and improve degraded rangelands after poor management. Successful reseeding will depend on; local climate, field characteristics, soil fertility, time of seeding, plant species selection, viability of seeds and the grazing management style.

#### 5. Shrub transplantation

To alleviate the spread of rangeland degradation, planting shrubs provides a large amount of fodder for livestock, combats desertification, and plays a key role in natural resource conservation. The integration of shrubs has the potential to improve both the sustainability and profitability of land, thus improving the livelihoods of smallholder farmers. Shrubs well-adapted to conditions of individual planting sites should be selected. The choice of species will depend on the annual rainfall rate, soil, topography, runoff, and water harvesting potential of the site, and the likelihood of environmental stresses such as drought, salinity, and cold. Species selection is also guided by rangeland development objectives, such as fodder production, wood production, dune fixation, or erosion control. On the other hand, the limits of this rehabilitation method can be translated by the high cost related to the nursery and transplantation; the necessity of protecting the site for more than 2-3 years to ensure plants establishment and survival. The use of exotic shrubs has shown discouraging results in most cases either due to their ecological demands or to the difficulty of their management.

#### 6. Grazing management

Grazing management is a tool to balance the capture of energy by the plants, the harvest of that energy by animals, and its conversion into a product that is marketable. Timing of grazing and growth rate of plants after grazing events are key factors in controlling the frequency, intensity and duration of grazing. These factors enhance soil stability, forage production, efficiency of forage use, and improve livestock production (Abu-Zanat *et al.*, 2005). Grazing is a natural method of land utilization to feed domestic livestock for converting grass and other forage into meat, milk and other products; it is the least expensive way to harvest forage. A proper grazing management should be holistic and using all the available resources with a pre-determined goal focusing on the fundamental ecological concepts/processes (Savory and Butterfield, 2016).

#### III - How does the toolbox work?

A toolbox is a guide that assists land managers and practitioners making diagnostic and taking proper measures to address causes of degradation. There is no "one size fits all", as every ecological site has its own specificity. Past disturbances have profound impact on the current degradation status. For instance, direct seeding in degraded rangelands needs to be combined with soil scarification, when the crust dominates the soil surface, to reactivate the soil water retention and increase seedling emergence and germination possibilities (Figure 1).




Fig. 1. Case study of the Jordanian Badia.

#### **Acknowledgments**

The authors would like to thank the Arab Fund for Social and Economic Development (AFSED) and the CGIAR Research Program (CRP) Livestock for their support and funding.

#### References

Abu-Zanat M.M., Migdady H.A. and Tabbaa M.J., 2005. Production systems of small ruminants in the Middle Badia of Jordan. *Dirasat, Agricultural Science* 32(2), 205-214.

Gauthier, M.M., Lambert, M.C. and Bédard, S., 2016. Effects of harvest gap size, soil scarification, and vegetation control on regeneration dynamics in sugar maple-yellow birch stands. *Forest Science*, 62(2), 237-246.

Milder J., Scherr S. and Bracer, C., 2010. Trends and future potential of payment for ecosystem services to alleviate rural poverty in developing countries. *Ecology and Society* 15(2), 4.

**Nefzaoui A., El Mourid M. and Louhaichi, M., 2014.** The Tribe – Platform of participatory local development and management of communal rangeland resources. *Journal of Arid Land Studies* 24(1), 57-60.

Ouled Belgacem A. and Louhaichi M., 2013. The vulnerability of native rangeland plant species to global climate change in the West Asia and North African regions. In: *Climatic Change* 119, 451-463.

- Oweis T.Y., 2016. Rainwater harvesting for restoring degraded dry agro-pastoral ecosystems: a conceptual review of opportunities and constraints in a changing climate. *Environmental reviews* 25(2), 135-149.
- Rist S., Chidambaranathan M., Escobar C., Wiesmann U. and Zimmermann A., 2007. Moving from sustainable management to sustainable governance of natural resources: the role of social learning processes in rural India, Bolivia and Mali. *Journal of Rural Studies* 23(1), 23-37.
- Savory A. and Butterfield J., 2016. Holistic management: A commonsense revolution to restore our environment. 3<sup>rd</sup> Edition, Island Press.

## **CIHEAM**

#### Centre International de Hautes Etudes Agronomiques Méditerranéennes

### **International Centre for Advanced Mediterranean Agronomic Studies**

Conseil d'Administration / Governing Board Président / President: Mohammed SADIKI

#### Délégués des Pays Membres / Member Countries Delegates

Albanie / Albania: Irfan TARELLI Liban / Lebanon: Wafaa DIKAH HAMZE

Algérie / Algeria: Farid HAROUADI Malte / Malta: Justin ZAHRA

Egypte / Egypt: Mohamed SOLIMAN Maroc / Morocco: Bilal HAJJOUJI

Espagne / Spain: María José HERNÁNDEZ MENDOZA Portugal: Nuno FIGUEIRA BOAVIDA CANADA

France: Valérie BADUEL Tunisia: Mahmoud Elies HAMZA

Grèce / Greece: Christos AVGOULAS Turquie / Turkey: Nihat PAKDIL

Italie / Italy: Teodoro MIANO

#### Comité Scientifique Consultatif / Scientific Advisory Committee

#### Président / President:

Mr Apostolos PAPADOPOULOS (Harokopio University of Athens, Greece)

#### **Membres / Members:**

Carole SINFORT (Montpellier SupAgro, France)

Lamiae GHAOUTI (Hassan II Institute of Agronomy and Veterinary Medicine, Morocco)

Giulio MALORGIO (University di Bologna, Italy)

Tomás GARCÍA AZCARATE (Institute of Economics, Geography and Demography, CSIC, Spain)

Maher Abdel-Mohsen ABDEL-HAMID (Cairo University, Faculty of Agriculture, Egypt)

**Suat SENSOY** (Yuzuncu Yil University, Turkey)

## **OPTIONS** méditerranéennes

**SERIES A: Mediterranean Seminars** 2021 – Number 125

#### Efficiency and resilience of forage resources and small ruminant production to cope with global challenges in Mediterranean areas

Edited by:

A. López-Francos, M. Jouven, C. Porqueddu, H. Ben Salem, A. Keli, A. Araba, M. Chentouf

The 1st Joint Meeting of the FAO-CIHEAM Network on Sheep and Goats (Subnetworks on Nutrition and Production Systems) and the FAO-CIHEAM Subnetwork on Mediterranean Pastures gathered more than 130 participants from 15 different countries in Meknes (Morocco), from 23 to 25 October 2019. The objective of the Meeting was to encourage the participation of and interaction between scientists, technicians and professionals to improve small ruminant productivity and to enhance the management of pastoral and forage resources in Mediterranean countries. The Meeting was devoted to present and exchange research advances on: (i) questions of common interest for the Networks (what production systems and value chains to meet the societal demand?, how to meet the challenge of climate change in Mediterranean agro-pastoral systems?), (ii) specific issues of interest for the Subnetworks (nutritional strategies to improve adaptation and production efficiency, promising forage resources for Mediterranean production systems); and (iii) the results from the 4-years research EU project iSAGE-Innovation for Sustainable Sheep and Goat production in Europe, dealt with in a specific Workshop. The Meeting was closed with a round table on pastoralists and extensive breeders, to present views and experiences in different countries to preserve and promote pastoral systems. This publication gathers the full articles of most of the presentations (keynotes, orals and posters) of the Meeting, providing a sample of the research efforts and scientific advances responding to the need to improve the efficiency and the resilience of Mediterranean forage resources and small ruminant production to cope with regional and global challenges.





ISBN: 978-2-85352-608-1

ISSN: 1016-121-X

## OPTIONS méditerranéennes

